

Generalized Master Function Approach to Quasi-Exactly Solvable Models

M. A. Jafarizadeh^{a,b,c} *, S. J. Akhtarshenas^{a,b,c} †

^aDepartment of Theoretical Physics and Astrophysics, Tabriz University, Tabriz 51664, Iran.

^bInstitute for Studies in Theoretical Physics and Mathematics, Tehran 19395-1795, Iran.

^cPure and Applied Science Research Center, Tabriz 51664, Iran.

February 7, 2008

*E-mail:jafarzadeh@ark.tabrizu.ac.ir

†E-mail:akhtarash@ark.tabrizu.ac.ir

Abstract

By introducing the generalized master function of order up to four together with corresponding weight function, we have obtained all quasi-exactly solvable second order differential equations. It is shown that these differential equations have solutions of polynomial type with factorization properties, that is polynomial solutions $P_m(E)$ can be factorized in terms of polynomial $P_{n+1}(E)$ for $m \geq n+1$. All known quasi-exactly quantum solvable models can be obtained from these differential equations, where roots of polynomial $P_{n+1}(E)$ are corresponding eigen-values.

Keywords: Quasi-Exact, Differential Equations, Orthogonal Polynomials, Factorization.

PACs Index: 03.65.Ge

1 INTRODUCTION

During the last decade a remarkable new class of quasi-exactly solvable spectral problems was introduced [1, 2, 3, 4, 5]. These occupy an intermediate position between exactly solvable and unsolvable models in the sense that exact solution in an algebraized form exists only for a part of the spectrum.

The usual approach to the analysis of quasi-exactly solvable systems is an algebraic one in which the operator is expressed as a non-linear combination of generators of a Lie algebra. Another recent developement is the work of Bender-Dunne [6] where they have shown that the eigen-functions of a quasi-exactly solvable schrodinger equation is the generating function for a set of orthogonal polynomials $P_m(E)$ in energy variable. It was further shown that, these polynomials satisfy the three-term recursion relation. Also, all polynomials beyond a critical polynomial $P_m(E)$ factorize into the product of polynomial $P_{n+1}(E)$ and another arbitrary polynomial.

In this paper we suggest a generalization of Bender-Dunne approach to all possible one-dimentional quasi-exactly second order differential equations.

For this purpose, the succesful master function approach of references [7, 8] to exactly solvable models, is generalized to a master function of order up to four which gives all possible one-dimensional quasi-exactly solvable models, where Bender-Dunne model [6] and Heun differential equation [9] are among them.

The paper is organized as follows: In section II we show that we can generalize the usual quadratic master function to a master function of at most four order polynomials, then the most general quasi-exactly solvable differential operators related to generalized master function of degree $k = 3$ and $k = 4$ are given, respectively.

In section III, expanding their solutions in powers of x , we get 3-term and 4-term recursion relations among their coefficeints, where Bender-Dunne factorization follows through imposing the quasi-exactly solvability conditions of section II. At the end of this section we list all possible related quasi-exactly solvable differential equations for $k = 3$ and $k = 4$ in Tables I and II, respectively.

Finally at section IV, we derive all possible one-dimensional quasi-exactly solvable quantum Hamiltonian

from the differential operators of section III, via prescription of references [7, 8], where we have listed them at the end of section III, except for those which can given in terms of elliptic functions. Paper ends with a brief conclusion.

2 QUASI-EXACTLY SOLVABLE DIFFERENTIAL EQUATIONS ASSOCIATED WITH GENERALIZED MASTER FUNCTIONS

By generalizing master function of order up to two [7, 8] to polynomial of order up to k , together with the non-negative weight function $W(x)$, defined at interval (a, b) such that $\frac{1}{W(x)} \frac{d}{dx} (A(x)W(x))$ to be a polynomial of degree at most $(k-1)$, we can define the operator

$$L = \frac{1}{W(x)} \frac{d}{dx} \left(A(x)W(x) \frac{d}{dx} \right) + B(x), \quad (2-1)$$

where $B(x)$ is a polynomial of order up to $(k-2)$. The interval (a, b) is chosen so that, we have $A(a)W(a) = A(b)W(b) = 0$.

It is straightforward to show that the above defined operator L is a self adjoint linear operator which at most maps a given polynomial of order m to another polynomial of order $(m+k-2)$. Now, by an appropriate choice of $B(x)$ and weight function $W(x)$, the operator L can have an invariant subspace of polynomials of order up to n . Then by choosing the set of orthogonal polynomials $\{\phi_0(x), \phi_1(x), \dots, \phi_n(x)\}$ defined in the interval (a, b) with respect to the weight function $W(x)$:

$$\int_a^b \phi_m(x) \phi_n(x) W(x) dx = 0, \quad \text{for } m \neq n \quad (2-2)$$

as the base, the matrix elements of the operator L on this base will have the following block diagonal form:

$$L_{ij} = 0, \quad \text{if } \{i \leq n \text{ and } j \geq n+1\} \text{ or } \{i \geq n+1 \text{ and } j \leq n\}. \quad (2-3)$$

Since, according to the well known theorem of orthogonal polynomials, $\phi_n(x)$ is orthogonal to any poly-

nomial of order up to $n - 1$, therefore, for matrix L we get

$$L = \begin{bmatrix} M & 0 \\ 0 & N \end{bmatrix}, \quad (2-4)$$

where M is an $(n + 1) \times (n + 1)$ matrix with matrix elements

$$M_{ij} = \int_a^b dx W(x) \phi_i(x) L(x) \phi_j(x), \quad i, j = 0, 1, 2, \dots, n, \quad (2-5)$$

and N is an infinite matrix element defined as above with $i, j \geq n + 1$.

The block diagonal form of the operator L indicates that by diagonalizing the $(n + 1) \times (n + 1)$ matrix M , we can find $(n + 1)$ eigen-values of the operator L together with the related eigen-functions as linear functions of orthogonal polynomials $\{\phi_0(x), \phi_1(x), \dots, \phi_n(x)\}$.

In order to determine the appropriate $B(x)$ and $W(x)$ for a given generalized master function $A(x)$, we Taylor expand those functions:

$$A(x) = \sum_{i=0}^k \frac{A^{(i)}(0)}{i!} x^i, \quad \text{where} \quad A^{(i)}(0) = \frac{d^i A(x)}{dx^i} \Big|_{x=0} \quad (2-6)$$

$$\frac{(A(x)W(x))'}{W(x)} = \sum_{i=0}^{k-1} \frac{\left(\frac{(AW)'}{W}\right)^{(i)}(0)}{i!} x^i, \quad \text{where} \quad \left(\frac{(AW)'}{W}\right)^{(i)}(0) = \frac{d^i \left(\frac{(A(x)W(x))'}{W(x)}\right)}{dx^i} \Big|_{x=0} \quad (2-7)$$

$$B(x) = \sum_{i=0}^{k-2} \frac{B^{(i)}(0)}{i!} x^i, \quad \text{where} \quad B^{(i)}(0) = \frac{d^i B(x)}{dx^i} \Big|_{x=0}. \quad (2-8)$$

Then, the existence of invariant subspace of the polynomials of order n of the operator L leads to the following linear equation between the coefficients of above Taylor expansions:

$$-\frac{A^{(i+2)}}{(i+2)!} l(l-1) - \frac{\left(\frac{(AW)'}{W}\right)^{(i+1)}}{(i+1)!} l + \frac{B^{(i)}}{i!} = 0, \quad (2-9)$$

where

$$\left\{ \begin{array}{l} l = n, \quad \text{and} \quad i = 1, \quad 2, \quad \dots, \quad k-2 \\ l = n-1, \quad \text{and} \quad i = 2, \quad 3, \quad \dots, \quad k-2 \\ \dots \quad \dots \quad \dots \quad \dots \quad \dots \quad \dots \\ l = n-k+4, \quad \text{and} \quad i = k-3, \quad k-2 \\ l = n-k+3, \quad \text{and} \quad i = k-2 \end{array} \right.$$

The number of above equations, for a given value of k , is $\frac{(k-1)(k-2)}{2}$. If we are to determine only the unknown function $B(x)$ without having any further constraint on the weight function $W(x)$, then the above $\frac{(k-1)(k-2)}{2}$ equations should be satisfied with $(k-2)$ coefficients of Taylor expansion of $B(x)$ as the only unknowns, since $B^{(0)}$ can be absorbed in the eigen-spectrum operator L . Therefore, we are left with $(k-2)$ unknowns to be determined, where the compatibility of equations (2-9) require $k = 3$ at most. On the other hand, if we add the coefficients of Taylor expansions of $A(x)$ and $\frac{(A(x)W(x))'}{W(x)}$ to our list of unknowns, (to be determined by solving equations (2-9)), then their compatibility conditions require that:

$$3(k-1) \geq \frac{(k-1)(k-2)}{2}, \quad (2-10)$$

or $k \leq 8$, where further investigations show that we can have at most $k = 4$, since for $k \geq 5$ the coefficients $A^{(k)}(0)$ and $\left(\frac{(AW)'}{W}\right)^{(k-1)}(0)$ will vanish. Below we summarize the above-mentioned discussion for $k = 3$ and $k = 4$, separately.

2.1 $k = 3$

In this case, $B(x)$ is a second order polynomial where $B^{(1)}$ can be determined by solving equation (2-9):

$$B^{(1)} = \frac{n}{2} \left(\frac{A^{(3)}}{3}(n-1) + \left(\frac{(AW)'}{W} \right)^{(2)} \right), \quad (2-11)$$

which is the only unknown in this case.

2.2 $k = 4$

Again, the solving of equations (2-9) leads to:

$$B^{(1)} = \frac{n}{2} \left(\frac{A^{(3)}}{3}(n-1) + \left(\frac{(AW)'}{W} \right)^{(2)} \right), \quad (2-12)$$

$$B^{(2)} = -\frac{A^{(4)}}{12}n(n-1), \quad (2-13)$$

and

$$\left(\frac{(AW)'}{W} \right)^{(3)} = -\frac{A^{(4)}}{2}(n-1). \quad (2-14)$$

Here, besides having constraint over second order polynomial $B(x)$, we have to put further constraints on the weight function $W(x)$ given in (2-14).

Definetly we can determine $n + 1$ eigen-spectrum of the operator L , simply by diagonalizing the $(n + 1) \times (n + 1)$ matrix M , since it is a self-adjoint operator in Hilbert space of polynomials and it has a block diagonal form given in (2-4).

As we are going to see in the next section, we can determine its eigen-spectrum analytically, using some recursion relations.

3 RECURSION RELATIONS

In this section we show that the eigen-functions of the operator L are a generating function for a new set of polynomials $P_m(E)$ where the eigen-function equation of the operator L leads to the recursion relation between these polynomials. Quasi-exact solvable constraints (2-9) will lead to their factorization, that is, $P_{n+N+1}(E) = P_{n+1}(E)Q_N(E)$ for $N \geq 0$, where roots of polynomials $P_{n+1}(E)$ turn out to be the eigen-values of the operator L .

To achieve these results, first we expand $\psi(x)$, the eigen-function of L , as:

$$\psi(x) = \sum_{m=0}^{\infty} P_m(E)x^m, \quad (3-15)$$

where eigen-function equation:

$$L\psi(x) = E\psi(x) \quad (3-16)$$

can be expressed as:

$$\begin{aligned} -A(x) \sum_{m=2}^{\infty} m(m-1)P_m(E)x^{m-2} - \frac{(A(x)W(x))'}{W(x)} \sum_{m=1}^{\infty} mP_m(E)x^{m-1} \\ + B(x) \sum_{m=0}^{\infty} P_m(E)x^m = E \sum_{m=0}^{\infty} P_m(E)x^m, \end{aligned} \quad (3-17)$$

which leads to the following recursion relations for the coefficients $P_m(E)$:

$$\begin{aligned} & \left(A^{(1)}(m+1)(m+2) + \left(\frac{(AW)'}{W} \right)^{(0)} (m+2) \right) P_{m+2}(E) \\ & + \left(\frac{A^{(2)}}{2!} m(m+1) + \left(\frac{(AW)'}{W} \right)^{(1)} (m+1) + E \right) P_{m+1}(E) \\ & + \left(\frac{A^{(3)}}{3!} m(m-1) + \frac{\left(\frac{(AW)'}{W} \right)^{(2)}}{2!} m - B^{(1)} \right) P_m(E) \\ & + \left(\frac{A^{(4)}}{4!} (m-1)(m-2) + \frac{\left(\frac{(AW)'}{W} \right)^{(3)}}{3!} m - \frac{B^{(2)}}{2!} \right) P_{m-1}(E) = 0. \end{aligned} \quad (3-18)$$

Below we investigate recursion relations thus obtained for $k = 3$ and $k = 4$, separately.

3.1 $k = 3$

In this case the 4-term general recursion relation reduce to the following 3-term recursion relation:

$$\begin{aligned} & \left(A^{(1)}(m+1)(m+2) + \left(\frac{(AW)'}{W} \right)^{(0)} (m+2) \right) P_{m+2}(E) \\ & + \left(\frac{A^{(2)}}{2!} m(m+1) + \left(\frac{(AW)'}{W} \right)^{(1)} (m+1) + E \right) P_{m+1}(E) \\ & + \left(\frac{A^{(3)}}{3!} m(m-1) + \frac{\left(\frac{(AW)'}{W} \right)^{(2)}}{2!} m - B^{(1)} \right) P_m(E) = 0. \end{aligned} \quad (3-19)$$

In order to have finite eigen-spectrum, that is, quasi-integrable differential equation, the above recursion relation should be truncated for some value of $m = n$, which is obviously possible by an appropriate choice of:

$$B^{(1)} = \frac{n}{2} \left(\frac{A^{(3)}}{3}(n-1) + \left(\frac{(AW)'}{W} \right)^{(2)} \right), \quad (3-20)$$

which is in agreement with the result of previous section given in (2-11).

Using the recursion relation (3-19), with $B^{(1)}$ given in (3-20), we get a factorization of polynomial $P_{n+N+1}(E)$ for $N \geq 0$ in terms of $P_{n+1}(E)$ as follows:

$$P_{n+N+1}(E) = P_{n+1}(E)Q_N(E) \quad N \geq 0 \quad (3-21)$$

where, by choosing the eigen-values E as roots of polynomial $P_{n+1}(E)$, all polynomials of order higher than n will vanish.

In order to determine corresponding eigen-functions, it is sufficient to evaluate $P_m(E_i)$ for $m = 0, 1, 2 \dots n$ with E_i as roots of $P_{n+1}(E)$, then eigen-function $\psi_i(x)$ corresponding to eigen-value E_i can be written as:

$$\psi_i(x) = \sum_{m=0}^n P_m(E_i)x^m, \quad i = 0, 1 \dots, n. \quad (3-22)$$

The above eigen-functions are polynomials of order n , hence they can have at most n roots in the interval (a, b) , where, according to the well known oscillation and comparison theorem of second-order linear differential equation [10], these numbers order the eigen-values according to the number of roots of corresponding eigen-functions. Therefore, we can say that the eigen-values thus obtained are the first $n + 1$ eigen-values of the operator L .

Using the recursion relations (3-19), we can evaluate the polynomials $P_m(E)$ in terms of $P_0(E)$, where we have chosen $P_0(E) = 1$. We have evaluated the first five polynomials which appear in the Appendix I.

As an illustration we give the results for $A(x) = x$ and $n = 3$ with $\alpha = 1, \beta = 0, \gamma = -1$ which is equivalent to the Bender-Dunne model:

$$P_1(E) = -\frac{1}{2}E,$$

$$P_2(E) = -1 + \frac{1}{12}E^2,$$

$$P_3(E) = \frac{1}{4}E - \frac{1}{144}E^3,$$

$$P_4(E) = \frac{1}{10} - \frac{1}{48}E^2 + \frac{1}{2880}E^4.$$

Obviously $P_m(E)$ have the parity of m .

By finding the 4-roots of $P_4(E)$ we determine the corresponding four eigen-values:

$$E_0 = -7.398556194,$$

$$E_1 = -2.293766823,$$

$$E_2 = 2.293766823,$$

$$E_3 = 7.398556194.$$

Finally for the coefficient $P_m(E_i)$ we get:

$$P_0(E_0) = 1, \quad P_1(E_0) = 3.699278097, \quad P_2(E_0) = 3.561552813, \quad P_3(E_0) = -.962769686,$$

$$P_0(E_1) = 1, \quad P_1(E_1) = 1.146883412, \quad P_2(E_1) = -.5615528135, \quad P_3(E_1) = -.4896337383,$$

$$P_0(E_2) = 1, \quad P_1(E_2) = -1.146883412, \quad P_2(E_2) = -.5615528135, \quad P_3(E_2) = .4896337383,$$

$$P_0(E_3) = 1, \quad P_1(E_3) = -3.699278097, \quad P_2(E_3) = 3.561552813, \quad P_3(E_3) = -.962769686.$$

Using the above coefficients we can determine the corresponding eigen-functions through formula (3-22).

In Table I we give all quasi-exactly solvable operators which can be obtained by choosing different generalized master function of order 3. This Table contains all possible models corresponding to different choice of $A(x)$ up to translation and rescaling of variable x . Also by choosing $A(x)$ as a polynomial of up to second order with $\gamma = 0$ we lead to the exactly solvable models of references [7, 8].

$A(x)$	$W(x)$	$L(x)$
	<i>intervals</i>	
x	$x^\alpha e^{\beta x + \gamma x^2}$ $0 \leq x < +\infty$ $\alpha > -1, -\infty < \beta < +\infty, \gamma < 0$	$-x \frac{d^2}{dx^2} - (\alpha + 1 + \beta x$ $+ 2\gamma x^2) \frac{d}{dx} + 2n\gamma x$
x^2	$x^\alpha e^{\beta/x + \gamma x}$ $0 \leq x < +\infty$ $-\infty < \alpha < +\infty, \beta < 0, \gamma < 0$	$-x^2 \frac{d^2}{dx^2} + (\beta - (\alpha + 2)x$ $-\gamma x^2) \frac{d}{dx} + n\gamma x$
$x(1-x)$	$x^\alpha (1-x)^\beta e^{-\gamma x}$ $0 \leq x \leq 1$ $\alpha > -1, \beta > -1, -\infty < \gamma < +\infty$	$x(x-1) \frac{d^2}{dx^2} + (-\alpha - 1$ $+ (\alpha + \beta + \gamma + 2)x$ $-\gamma x^2) \frac{d}{dx} + n\gamma x$
x^3	$x^\alpha e^{-\beta/x^2 - \gamma/x}$ $0 \leq x < +\infty$ $\alpha < -3, \beta > 0, -\infty < \gamma < +\infty$	$-x^3 \frac{d^2}{dx^2} - (2\beta + \gamma x$ $+ (\alpha + 3)x^2) \frac{d}{dx} + n(n + \alpha + 2)x$
$x^2(1-x)$	$x^\alpha (1-x)^\beta e^{-\gamma/x}$ $0 \leq x \leq 1$ $-\infty < \alpha < +\infty, \beta > -1, \gamma > 0$	$x^2(x-1) \frac{d^2}{dx^2} + (-\gamma + (\gamma - \alpha - 2)x$ $+ (\alpha + \beta + 3)x^2) \frac{d}{dx}$ $-n(n + \alpha + \beta + 2)x$
$x(1+x^2)$	$x^\alpha (1+x^2)^\beta e^{\gamma \tan^{-1} x}$ $0 \leq x < +\infty$ $\alpha > -1, \beta < -(\alpha + 3)/2, -\infty < \gamma < +\infty$	$-x(1+x^2) \frac{d^2}{dx^2} - (\alpha + 1 + \gamma x$ $+ (\alpha + 2\beta + 3)x^2) \frac{d}{dx}$ $+ n(n + \alpha + 2\beta + 2)x$
$x(1-x)(a-x)$ $a > 1$	$x^\alpha (1-x)^\beta (a-x)^\gamma$ $0 \leq x \leq 1$ $\alpha > -1, \beta > -1, -\infty < \gamma < +\infty$	$x(x-1)(a-x) \frac{d^2}{dx^2} + (-a(\alpha + 1)$ $+ ((a+1)(\alpha + 2) + a\beta + \gamma)x - (\alpha + \beta$ $+ \gamma + 3)x^2) \frac{d}{dx} + n(n + \alpha + \beta + \gamma + 2)x$

Table 1: Quasi-exactly differential operators obtained from generalized master function of order up to 3

3.2 $k = 4$

Again in order to truncate the recursion relation (3-18) and to factorize polynomials $P_{n+N+1}(E)$ in terms of $P_{n+1}(E)$, we should have:

$$B^{(1)} = \frac{n}{2} \left(\frac{A^{(3)}}{3}(n-1) + \left(\frac{(AW)'}{W} \right)^{(2)} \right), \quad (3-23)$$

$$\frac{B^{(2)}}{2!} = \frac{A^{(4)}}{4!}(n-1)(n-2) + \frac{\left(\frac{(AW)'}{W} \right)^{(3)}}{3!}n, \quad (3-24)$$

and

$$\frac{B^{(2)}}{2!} = \frac{A^{(4)}}{4!}n(n-1) + \frac{\left(\frac{(AW)'}{W} \right)^{(3)}}{3!}(n+1). \quad (3-25)$$

Solving the above equations we get:

$$B^{(2)} = -\frac{A^{(4)}}{12}n(n-1), \quad (3-26)$$

and

$$\left(\frac{(AW)'}{W} \right)^{(3)} = -\frac{A^{(4)}}{2}(n-1). \quad (3-27)$$

The equations (3-23) , (3-26) and (3-27) are the same equations which are required in the reduction of the operator L to its block diagonal form.

Again roots of polynomials $P_{n+1}(E)$ will correspond to $n+1$ eigen-values of the differential operator L with eigen-functions which can be expressed in terms of $P_m(E_i)$ for $m \leq n$, where polynomials $P_m(E)$ can be obtained from recursion relation by choosing $P_0(E) = 1$ and $P_{-1}(E) = 0$, where we have given the first four polynomials in Appendix II.

In Table II we list all quasi-exactly differential operators which can be obtained from the generalized master function of order up to four.

Tables I and II contain all quasi-exactly second order differential equations which can be obtained from Lie algebraic methods [4]. For example we get the Bender-Dunne model [6] for the choice of $A(x) = x$ and $\beta = 0$, and similarly, we get the Heun differential operator (Fuxian equation with four regular singular point) [9], for the choice of $A(x) = x(x-1)(x-a)$.

$A(x)$	$W(x)$	$L(x)$
	<i>intervals</i>	
x^4	$x^\alpha e^{\beta/x^3 + \gamma/x^2 + \delta/x}$ $0 \leq x < +\infty$ $\beta < 0, -\infty < \gamma < +\infty, -\infty < \delta < +\infty$ $\alpha = -2(n+1)$	$-x^4 \frac{d^2}{dx^2} + (3\beta + 2\gamma x$ $+ \delta x^2 - (\alpha + 4)x^3) \frac{d}{dx} - n\delta x - n(n-1)x^2$
$x^3(1-x)$	$x^\alpha (1-x)^\beta e^{-\gamma/x - \delta/x^2}$ $0 \leq x \leq +1$ $\beta > -1, -\infty < \gamma < +\infty, \delta > 0$ $\alpha = -2(n+1) - \beta$	$x^3(x-1) \frac{d^2}{dx^2} + (-2\delta + (2\delta - \gamma)x$ $+ (\gamma - \alpha - 3)x^2 + (\alpha + \beta + 4)x^3) \frac{d}{dx}$ $+ n(n + \alpha - \gamma + 2)x + n(n-1)x^2$
$x^2(1+x^2)$	$x^\alpha (1+x^2)^\beta e^{\gamma/x + \delta \tan^{-1}(x)}$ $0 \leq x < +\infty$ $-\infty < \beta < +\infty, \gamma < 0, -\infty < \delta < +\infty$ $\alpha = -2(n + \beta + 1)$	$-x^2(1+x^2) \frac{d^2}{dx^2} + (\gamma - (\alpha + 2)x$ $+ (\gamma - \delta)x^2 - (\alpha + 2\beta + 4)x^3) \frac{d}{dx}$ $+ n(\delta - \gamma)x - n(n-1)x^2$
$x^2(1-x)(a-x)$ $a > 1$	$x^\alpha (1-x)^\beta (a-x)^\gamma e^{\delta/x}$ $0 \leq x \leq +1$ $\beta > -1, -\infty < \gamma < +\infty, \delta < 0$ $\alpha = -2(n+1) - \beta - \gamma$	$x^2(x-1)(a-x) \frac{d^2}{dx^2} + (a\delta - (a\alpha + (a+1)\delta$ $+ 2a)x - ((a+1)\alpha + a\beta + \gamma + \delta$ $+ 3(a+1))x^2 - (\alpha + \beta + \gamma + 4)x^3) \frac{d}{dx} + n((\alpha$ $- n + 4)(a+1) + a\beta + \gamma + \delta)x - n(n-1)x^2$
$x^2(1-x)^2$	$x^\alpha (1-x)^\beta e^{\gamma/x + \delta/(1-x)}$ $0 \leq x \leq +1$ $-\infty < \beta < +\infty, \gamma < 0, \delta < 0$ $\alpha = -2(n+1) - \beta$	$-x^2(1-x)^2 \frac{d^2}{dx^2} + (\gamma - (\alpha + 2\gamma + 2)x$ $+ (2\alpha + \beta + \gamma - \delta + 6)x^2 - (\alpha + \beta + 4)x^3) \frac{d}{dx}$ $- n(+2n + 2\alpha + \beta + \gamma - \delta + 4)x - n(n-1)x^2$
$x(a-x)(1+x^2)$ $a > 0$	$x^\alpha (a-x)^\beta (1+x^2)^\gamma e^{\delta \tan^{-1}x}$ $0 \leq x \leq a$ $-\infty < \gamma < +\infty, -\infty < \delta < +\infty$ $-1 < \beta < -2n - 2\gamma - 1$ $\alpha = -2(n+1) - \beta - 2\gamma$	$x(x-a)(1+x^2) \frac{d^2}{dx^2} + (-a(\alpha + 1)$ $+ (\alpha + \beta - a(\delta - 2))x - (a(\alpha + 2\gamma + 3)$ $- \delta)x^2 + (\alpha + \beta + 2\gamma + 4)x^3) \frac{d}{dx}$ $+ n(a(n + \alpha + 2\gamma + 2) - \delta)x + n(n-1)x^2$
$x(1-x)(a-x)(b-x)$ $1 < a < b$	$x^\alpha (1-x)^\beta (a-x)^\gamma (b-x)^\delta$ $0 \leq x \leq +1$ $-1 < \beta < -2n - \gamma - 1$	$x(x-1)(a-x)(b-x) \frac{d^2}{dx^2}$ $+ (-ab(\alpha + 1) + (2(a+b+ab) + ab(\alpha + \beta)$ $+ b(\alpha + \gamma) + a(\alpha + \delta))x - ((a+b)(\alpha + \beta + 3)$

4 QUASI-EXACTLY POTENTIAL ASSOCIATED WITH GENERALIZED MASTER FUNCTION

As in references [7, 8], writing:

$$\psi(t) = A^{1/4}(x)W^{1/2}(x)\phi(x), \quad (4-28)$$

with a change of variable $\frac{dx}{dt} = \sqrt{A(x)}$, the eigen-value equation of the operator L reduces to the schrodinger equation:

$$H(t)\psi(t) = E\psi(t), \quad (4-29)$$

with the same eigen-value E and $\psi(t)$ given in (4-29), in terms of the eigen function of L , where $H(t) = -\frac{d^2}{dt^2} + V(t)$ is the similarity transformation of $L(x)$ defined as:

$$H(t) = A^{1/4}(x)W^{1/2}(x)L(x)A^{-1/4}(x)W^{-1/2}(x) \quad (4-30)$$

with:

$$V(t) = -\frac{3}{16} \frac{\dot{A}^2(t)}{A^2(t)} - \frac{1}{4} \frac{\dot{W}^2(t)}{W^2(t)} + \frac{1}{4} \frac{\dot{A}(t)\dot{W}(t)}{A(t)W(t)} + \frac{1}{4} \frac{\ddot{A}(t)}{A(t)} + \frac{1}{2} \frac{\ddot{W}(t)}{W(t)} + B(t). \quad (4-31)$$

It is also straightforward to show that:

$$\int dt \phi(t)H(t)\psi(t) = \int_a^b dx W(x)\psi(x)L(x)\psi(x). \quad (4-32)$$

Hence block diagonalization of L leads to block-diagonalization of H .

As an illustration we give below an example with $A(x) = x^3$, weight function $W(x) = x^\alpha e^{\beta/x^2 - \gamma/x}$ and interval $[0, \infty)$, where $\alpha < -3$ and $\beta, \gamma > 0$.

From a change of variable $dx/dt = \sqrt{A(x)}$, we get $x(t) = 4/t^2$, hence using equation (4-31) we have for potential $V(t)$:

$$\begin{aligned} V_n(t) = & \frac{\gamma}{2}(\alpha + 1) + \left(\frac{15}{4} + \alpha^2 + 4n\alpha + 4\alpha + 4n^2 + 8n \right) \frac{1}{t^2} + \frac{1}{4} \left(\alpha\beta + \frac{1}{4}\gamma^2 \right) t^2 \\ & + \frac{\beta\gamma}{16} t^4 + \frac{\beta^2}{64} t^6 \end{aligned} \quad (4-33)$$

Below we give a list of quasi-exact solvable potentials, except for those potentials which can be expressed in terms of elliptic functions, since in this case we get rather long expressions for them:

$$A(x) = x, \quad x(t) = \frac{t^2}{4}$$

$$V_n(t) = \frac{\beta}{2}(\alpha + 1) + \left(\alpha^2 - \frac{1}{4}\right) \frac{1}{t^2} + \frac{1}{2} \left(\frac{\beta^2}{8} + \gamma \left(n + 1 + \frac{\alpha}{2}\right)\right) t^2 + \frac{\beta\gamma}{16} t^4 + \frac{\gamma^2}{64} t^6,$$

$$A(x) = x^2, \quad x(t) = e^t$$

$$V_n(t) = \frac{1}{4} \left(1 + \alpha^2 - 2\beta\gamma + 2\alpha - 2\alpha\beta e^{-t} + \beta^2 e^{-2t} + \gamma^2 e^{2t} + 2(2\gamma + 2n\gamma + \alpha\gamma) e^t\right),$$

$$A(x) = x(1 - x), \quad x(t) = \frac{1 + \sin(t)}{2}$$

$$V_n(t) = \frac{1}{2} \left(+n\gamma - \alpha\beta - \beta - \alpha + \frac{1}{2}(\beta\gamma - \alpha^2 - \beta^2 - \alpha\gamma - 1) + \left(\frac{\alpha\gamma}{2} + \gamma + \frac{\beta\gamma}{2} + n\gamma\right) \sin(t) \right) \\ \frac{1}{2} \left(\alpha^2 + \beta^2 - \frac{1}{2} + (\beta^2 - \alpha^2) \sin(t) \right) \frac{1}{\cos^2(t)} + \frac{\gamma^2}{16} \cos^2(t),$$

$$A(x) = x^3, \quad x(t) = \frac{4}{t^2}$$

$$V_n(t) = \frac{\gamma}{2}(\alpha + 1) + \left(\frac{15}{4} + \alpha^2 + 4n\alpha + 4\alpha + 4n^2 + 8n\right) \frac{1}{t^2} + \frac{1}{4} \left(\alpha\beta + \frac{\gamma^2}{4}\right) t^2 \\ + \frac{\beta\gamma}{16} t^4 + \frac{\beta^2}{64} t^6,$$

$$A(x) = x^2(1 - x), \quad x(t) = 1 - \tanh^2\left(\frac{t}{2}\right)$$

$$V_n(t) = \left(\frac{1}{\cosh^2(t) - 1}\right) \left(-\left(2n^2 + 2 + 2n\alpha + \frac{\alpha^2}{2} + 4n + \alpha\beta + 2\alpha + \frac{\alpha\gamma}{4} + 2n\beta + 2\beta\right) \cosh(t) \right. \\ \left. + \frac{1}{2} \left(\frac{-\gamma^2}{4} - \frac{\alpha\gamma}{2} + \frac{1}{2} - \gamma + \alpha - \beta\gamma + \frac{\alpha^2}{2}\right) \cosh^2 + \frac{\alpha\gamma}{4} \cosh^3 + \frac{\gamma^2}{16} \cosh^4\right)$$

$$+ \left(\frac{\gamma}{2} + 4n + \alpha\beta + \frac{3\alpha}{2} + \beta^2 + 2\beta + \frac{3}{2} + \frac{\alpha^2}{4} + \frac{\alpha\gamma}{4} + \frac{\beta\gamma}{2} + \frac{\gamma^2}{16} + 2n\beta + 2n^2 + 2n\alpha \right),$$

$$A(x) = x^4, \quad x(t) = \frac{1}{t}$$

$$V_n(t) = \left(\frac{\delta^2}{4} + \gamma + 2n\gamma \right) + (\gamma\delta + 3n\beta + 3\beta)t + \left(\frac{3\beta\delta}{2} + \gamma^2 \right)t^2 + 3\beta\gamma t^3 + \frac{9\beta^2}{4}t^4,$$

$$A(x) = x^3(1-x), \quad x(t) = \frac{4}{4+t^2}$$

$$V_n(t) = - \left(\frac{\gamma}{2} + \delta + \beta\delta + \frac{\beta\gamma}{2} + n\gamma \right) + \left(\beta^2 - \frac{1}{4} \right) \frac{1}{t^2} \\ + \frac{1}{2} \left(-n\delta + \frac{\delta^2}{2} - \frac{\beta\delta}{2} - \delta + \frac{\gamma^2}{8} + \frac{\gamma\delta}{2} \right) t^2 + \frac{\delta}{8} \left(\frac{\gamma}{2} + \delta \right) t^4 + \frac{\delta^2}{64} t^6,$$

$$A(x) = x^2(1+x^2), \quad x(t) = \frac{-1}{\sinh(t)}$$

$$V_n(t) = \left(n + n^2 - \frac{\gamma\delta}{2} + \frac{1}{4} + 2n\beta + \beta + \beta^2 - (n\gamma + \gamma + \beta\gamma) \sinh(t) \right) \\ + \left(\frac{\delta^2}{4} + \beta\delta \sinh(t) - \beta^2 + \frac{1}{4} \right) \frac{1}{\cosh^2(t)} + \frac{\gamma^2}{4} \cosh^2(t),$$

$$A(x) = x^2(1-x^2), \quad x(t) = \frac{1}{\cosh(t)}$$

$$V_n(t) = \left(\frac{1}{\cosh^2(t) - 1} \right) \left(\left(\frac{-\gamma}{2} - n - \frac{\beta}{2} + \frac{\gamma^2}{4} + \frac{\delta^2}{4} - \frac{\beta\gamma}{2} - \frac{\beta\delta}{2} + \frac{\gamma\delta}{2} - \frac{1}{2} - n\beta + \frac{\beta^2}{4} - n\gamma - n^2 \right) \right. \\ + \left(\frac{\beta\delta}{2} + n\beta - \frac{\gamma\delta}{2} + n\gamma + \frac{\gamma}{2} + \frac{1}{4} + \frac{\beta}{2} + \frac{\beta\gamma}{2} - \frac{\delta^2}{2} + n^2 + \frac{\beta^2}{4} + n + \frac{\gamma^2}{4} \right) \cosh(t)^2 \\ \left. + \left(\frac{-\gamma\delta}{2} - n\delta - \delta - \frac{\gamma^2}{2} - \frac{\beta\delta}{2} + \frac{\beta^2}{2} \right) \cosh(t) + \left(\delta + n\delta + \frac{\beta\delta}{2} + \frac{\gamma\delta}{2} \right) \cosh^3(t) + \frac{\delta^2}{4} \cosh^4(t) \right),$$

$$A(x) = x^2(1-x)^2, \quad x(t) = \frac{e^t}{1+e^t}$$

$$\begin{aligned}
V_n(t) = & \left(\frac{1}{(e^{-t} + 1)^4} S_6 \right) \left(\frac{\delta^2}{4} S_4 + \delta \left(\delta - \frac{\beta}{2} \right) S_5 \right. \\
& + \left(\frac{1}{4} - \frac{\gamma\delta}{2} + n^2 + \frac{\beta^2}{4} - n\gamma + \frac{\beta}{2} + n - 2\beta\delta + \frac{3\delta^2}{2} + n\beta \right) S_6 \\
& + \left(-3\beta\delta + \frac{\beta\gamma}{2} - 3n\gamma + \gamma + 4n + \delta^2 - 2\gamma\delta + \beta^2 + 1 + 4n^2 + 2\beta + 4n\beta \right) S_7 \\
& + \left(3\beta + 6n + \frac{3\beta^2}{2} - 3\gamma\delta + \frac{3}{2} + \frac{\delta^2}{4} + \frac{\gamma^2}{4} + 6n^2 - 2\beta\delta + 4\gamma + 6n\beta + 2\beta\gamma - 2n\gamma \right) S_8 \\
& + \left(4n^2 - \frac{\beta\delta}{2} - 2\gamma\delta + 3\beta\gamma + 1 + 2n\gamma + 4n + \beta^2 + 2\beta + \gamma^2 + 6\gamma + 4n\beta \right) S_9 \\
& + \left(\frac{-\gamma\delta}{2} + \frac{3\gamma^2}{2} + 4\gamma + n + 2\beta\gamma + 3n\gamma + \frac{1}{4} + \frac{\beta^2}{4} + \frac{\beta}{2} + n\beta + n^2 \right) S_{10} \\
& \left. + \left(n\gamma + \gamma^2 + \frac{\beta\gamma}{2} + \gamma \right) S_{11} + \frac{\gamma^2}{4} S_{12} \right),
\end{aligned}$$

where S_k is defined as:

$$S_k = \exp \left(-2 \left(\frac{k}{2} t + (\gamma e^{-t} + \gamma + \delta (e^t + 1)) \right) \right), \quad k = 4, 5, \dots, 12.$$

5 CONCLUSION

As we saw by introducing of master function $A(x)$ as a polynomial of order at most four, we could obtain all quasi-exactly second order differential equations. It is shown that the eigen-equation relation $L\Psi(x) = E\Psi(x)$ generates a set of polynomials $P_m(E)$, where these polynomials satisfy 3-term and 4-term recursion relations for master function of at most three and four, respectively. Finally the quasi-exactly solvability leads to factorization of polynomials $P_{n+N+1}(E)$ for $N \geq 0$ in terms of $P_{n+1}(E)$, where by determining the roots of $P_{n+1}(E)$ we can determine first $n + 1$ eigen-values of these quasi-exactly solvable differential equations.

APPENDIX I

The First Five Polynomials $P_n(E)$, For $k = 3$

To abbreviate, we set $F^{(i)} = \left(\frac{AW'}{W} \right)^{(i)}$,

$$P_1(E) = -\frac{E}{F^{(0)}},$$

$$P_2(E) = \frac{1}{2} \frac{EF^{(1)} + E^2 + B^{(1)} F^{(0)}}{F^{(0)} (A^{(1)} + F^{(0)})},$$

$$P_3(E) = \left(-A^{(2)} EF^{(1)} - A^{(2)} E^2 - A^{(2)} B^{(1)} F^{(0)} - 2EF^{(1)^2} - 3F^{(1)} E^2 - 2F^{(1)} B^{(1)} F^{(0)} - E^3 \right. \\ \left. - 3EB^{(1)} F^{(0)} + EF^{(2)} A^{(1)} + EF^{(2)} F^{(0)} - 2EB^{(1)} A^{(1)} \right) \\ \left/ \left(6F^{(0)} (A^{(1)} + F^{(0)}) (2A^{(1)} + F^{(0)}) \right) \right.,$$

$$P_4(E) = - \left(-3A^{(2)^2} E^2 - 3B^{(1)^2} F^{(0)^2} - 6B^{(1)^2} F^{(0)} A^{(1)} + 3F^{(2)} B^{(1)} F^{(0)^2} + A^{(3)} B^{(1)} F^{(0)^2} \right. \\ \left. + A^{(3)} E^2 F^{(0)} + 2A^{(3)} E^2 A^{(1)} - 8E^2 B^{(1)} A^{(1)} + 4E^2 F^{(2)} F^{(0)} + 7E^2 F^{(2)} A^{(1)} - 6E^2 B^{(1)} F^{(0)} \right. \\ \left. - 6F^{(1)^2} B^{(1)} F^{(0)} - 13A^{(2)} F^{(1)} E^2 - 9A^{(2)} EF^{(1)^2} - 3A^{(2)^2} B^{(1)} F^{(0)} - 3A^{(2)^2} EF^{(1)} \right. \\ \left. - 4A^{(2)} E^3 - 6EF^{(1)^3} - 11F^{(1)^2} E^2 - 6F^{(1)} E^3 - 9A^{(2)} F^{(1)} B^{(1)} F^{(0)} - 6A^{(2)} EB^{(1)} A^{(1)} \right. \\ \left. + 3A^{(2)} EF^{(2)} F^{(0)} + 3A^{(2)} EF^{(2)} A^{(1)} - 10A^{(2)} EB^{(1)} F^{(0)} - 12F^{(1)} EB^{(1)} A^{(1)} + 6F^{(1)} EF^{(2)} F^{(0)} \right. \\ \left. + 9F^{(1)} EF^{(2)} A^{(1)} - 14F^{(1)} EB^{(1)} F^{(0)} + 2A^{(3)} B^{(1)} F^{(0)} A^{(1)} + A^{(3)} EF^{(1)} F^{(0)} \right. \\ \left. + 2A^{(3)} EF^{(1)} A^{(1)} + 6F^{(2)} B^{(1)} F^{(0)} A^{(1)} - E^4 \right) \\ \left/ \left(24F^{(0)} (A^{(1)} + F^{(0)}) (2A^{(1)} + F^{(0)}) (3A^{(1)} + F^{(0)}) \right) \right.,$$

$$\begin{aligned}
P_5(E) = & - \left(-46 E A^{(3)} B^{(1)} F^{(0)} A^{(1)} - 88 E F^{(2)} B^{(1)} F^{(0)} A^{(1)} + 16 A^{(3)} E F^{(2)} A^{(1)} F^{(0)} \right. \\
& + 24 E F^{(2)^2} A^{(1)} F^{(0)} - 48 F^{(2)} E B^{(1)} A^{(1)^2} + 18 E F^{(2)^2} A^{(1)^2} + 6 E F^{(2)^2} F^{(0)^2} + 24 E B^{(1)^2} A^{(1)^2} \\
& + 12 A^{(3)} E F^{(2)} A^{(1)^2} + 4 A^{(3)} E F^{(2)} F^{(0)^2} - 24 A^{(3)} E B^{(1)} A^{(1)^2} - 32 F^{(1)} A^{(3)} B^{(1)} F^{(0)} A^{(1)} \\
& - 60 F^{(1)} F^{(2)} B^{(1)} F^{(0)} A^{(1)} + 50 E B^{(1)^2} F^{(0)} A^{(1)} - 25 E F^{(2)} B^{(1)} F^{(0)^2} - 13 E A^{(3)} B^{(1)} F^{(0)^2} \\
& - 46 F^{(1)} A^{(3)} E^2 A^{(1)} + 80 F^{(1)} E^2 B^{(1)} A^{(1)} - 40 F^{(1)} E^2 F^{(2)} F^{(0)} - 91 F^{(1)} E^2 F^{(2)} A^{(1)} \\
& + 50 F^{(1)} E^2 B^{(1)} F^{(0)} - 54 A^{(2)} F^{(1)} E F^{(2)} F^{(0)} - 84 A^{(2)} F^{(1)} E F^{(2)} A^{(1)} + 137 A^{(2)} F^{(1)} E B^{(1)} F^{(0)} \\
& - 24 A^{(2)} A^{(3)} B^{(1)} F^{(0)} A^{(1)} - 10 A^{(2)} A^{(3)} E F^{(1)} F^{(0)} - 24 A^{(2)} A^{(3)} E F^{(1)} A^{(1)} \\
& - 54 A^{(2)} F^{(2)} B^{(1)} F^{(0)} A^{(1)} + 20 F^{(1)} B^{(1)^2} F^{(0)^2} - 5 A^{(3)} E^3 F^{(0)} - 14 A^{(3)} E^3 A^{(1)} + 20 E^3 B^{(1)} A^{(1)} \\
& - 10 E^3 F^{(2)} F^{(0)} - 25 E^3 F^{(2)} A^{(1)} + 10 E^3 B^{(1)} F^{(0)} + 15 E B^{(1)^2} F^{(0)^2} + 66 A^{(2)} E^2 B^{(1)} A^{(1)} \\
& - 33 A^{(2)} E^2 F^{(2)} F^{(0)} - 63 A^{(2)} E^2 F^{(2)} A^{(1)} + 50 A^{(2)} E^2 B^{(1)} F^{(0)} + 72 A^{(2)} F^{(1)^2} B^{(1)} F^{(0)} \\
& + 72 F^{(1)^2} E B^{(1)} A^{(1)} - 36 F^{(1)^2} E F^{(2)} F^{(0)} - 72 F^{(1)^2} E F^{(2)} A^{(1)} + 70 F^{(1)^2} E B^{(1)} F^{(0)} \\
& - 12 A^{(3)} E F^{(1)^2} F^{(0)} - 32 A^{(3)} E F^{(1)^2} A^{(1)} + 48 F^{(1)} B^{(1)^2} F^{(0)} A^{(1)} - 24 F^{(1)} F^{(2)} B^{(1)} F^{(0)^2} \\
& - 12 F^{(1)} A^{(3)} B^{(1)} F^{(0)^2} - 17 F^{(1)} A^{(3)} E^2 F^{(0)} - 24 A^{(2)} F^{(2)} B^{(1)} F^{(0)^2} - 10 A^{(2)} A^{(3)} E^2 F^{(0)} \\
& - 24 A^{(2)} A^{(3)} E^2 A^{(1)} - 10 A^{(2)} A^{(3)} B^{(1)} F^{(0)^2} + 108 A^{(2)} F^{(1)} E B^{(1)} A^{(1)} \\
& + 10 F^{(1)} E^4 + 18 A^{(2)^3} E^2 + 27 A^{(2)^2} E^3 + 10 A^{(2)} E^4 + 24 E F^{(1)^4} + 50 F^{(1)^3} E^2 \\
& + 35 F^{(1)^2} E^3 + E^5 + 93 A^{(2)^2} F^{(1)} E^2 + 66 A^{(2)^2} E F^{(1)^2} + 18 A^{(2)^3} B^{(1)} F^{(0)} + 18 A^{(2)^3} E F^{(1)} \\
& + 22 A^{(2)} B^{(1)^2} F^{(0)^2} + 72 A^{(2)} E F^{(1)^3} + 127 A^{(2)} F^{(1)^2} E^2 + 65 A^{(2)} F^{(1)} E^3 \\
& + 24 F^{(1)^3} B^{(1)} F^{(0)} + 66 A^{(2)^2} F^{(1)} B^{(1)} F^{(0)} + 36 A^{(2)^2} E B^{(1)} A^{(1)} \\
& - 18 A^{(2)^2} E F^{(2)} F^{(0)} - 18 A^{(2)^2} E F^{(2)} A^{(1)} + 63 A^{(2)^2} E B^{(1)} F^{(0)} + 48 A^{(2)} B^{(1)^2} F^{(0)} A^{(1)} \Big) \\
& \Big/ \left(120 F^{(0)} \left(A^{(1)} + F^{(0)} \right) \left(2 A^{(1)} + F^{(0)} \right) \left(3 A^{(1)} + F^{(0)} \right) \left(4 A^{(1)} + F^{(0)} \right) \right).
\end{aligned}$$

APPENDIX II

The First Four Polynomials $P_n(E)$, For $k = 4$

$$P_1(E) = \frac{E}{F^{(0)}},$$

$$P_2(E) = -\frac{1}{2} \frac{EF^{(1)} + E^2 - B^{(1)} F^{(0)}}{F^{(0)} (A^{(1)} + F^{(0)})},$$

$$P_3(E) = \left(A^{(2)} EF^{(1)} + A^{(2)} E^2 - A^{(2)} B^{(1)} F^{(0)} + 2 EF^{(1)2} + 3 F^{(1)} E^2 - 2 F^{(1)} B^{(1)} F^{(0)} \right. \\ \left. + E^3 + EB^{(1)} F^{(0)} - EF^{(2)} A^{(1)} - EF^{(2)} F^{(0)} + 2 EB^{(1)} A^{(1)} + B^{(2)} F^{(0)} A^{(1)} + B^{(2)} F^{(0)2} \right) \\ \left/ \left(6F^{(0)} (A^{(1)} + F^{(0)}) (2A^{(1)} + F^{(0)}) \right) \right.,$$

$$P_4(E) = - \left(3 A^{(2)} B^{(2)} F^{(0)2} + A^{(3)} B^{(1)} F^{(0)2} + 9 A^{(2)} EF^{(1)2} - 3 A^{(2)} EF^{(2)} A^{(1)} \right. \\ \left. + 2 A^{(2)} EB^{(1)} F^{(0)} - 9 A^{(2)} F^{(1)} B^{(1)} F^{(0)} - 2 EB^{(2)} F^{(0)2} + 3 A^{(2)} B^{(2)} F^{(0)} A^{(1)} + 6 A^{(2)} EB^{(1)} A^{(1)} \right. \\ \left. - 3 A^{(2)} EF^{(2)} F^{(0)} - 9 F^{(1)} EF^{(2)} A^{(1)} + 4 F^{(1)} EB^{(1)} F^{(0)} - 3 B^{(1)2} F^{(0)2} - 8 EB^{(2)} F^{(0)} A^{(1)} \right. \\ \left. - 2 A^{(3)} EF^{(1)} A^{(1)} - A^{(3)} EF^{(1)} F^{(0)} + 2 A^{(3)} B^{(1)} F^{(0)} A^{(1)} + 6 F^{(2)} B^{(1)} F^{(0)} A^{(1)} + 3 EF^{(3)} A^{(1)} F^{(0)} \right. \\ \left. - 6 F^{(1)} EF^{(2)} F^{(0)} + 3 A^{(2)2} E^2 + 4 A^{(2)} E^3 + 6 EF^{(1)3} + 11 F^{(1)2} E^2 + 6 F^{(1)} E^3 + 3 A^{(2)2} EF^{(1)} \right. \\ \left. - 3 A^{(2)2} B^{(1)} F^{(0)} + 13 A^{(2)} F^{(1)} E^2 - 6 F^{(1)2} B^{(1)} F^{(0)} + 3 F^{(1)} B^{(2)} F^{(0)2} + 4 E^2 B^{(1)} F^{(0)} \right. \\ \left. - 7 E^2 F^{(2)} A^{(1)} - 4 E^2 F^{(2)} F^{(0)} + 8 E^2 B^{(1)} A^{(1)} - 2 A^{(3)} E^2 A^{(1)} - A^{(3)} E^2 F^{(0)} + 3 F^{(2)} B^{(1)} F^{(0)2} \right. \\ \left. - 6 B^{(1)2} F^{(0)} A^{(1)} + 2 EF^{(3)} A^{(1)2} + EF^{(3)} F^{(0)2} - 6 EB^{(2)} A^{(1)2} + E^4 + 3 F^{(1)} B^{(2)} F^{(0)} A^{(1)} \right)$$

$$+12 F^{(1)} E B^{(1)} A^{(1)} \Big) \Big/ \Big(24 F^{(0)} \Big(A^{(1)} + F^{(0)} \Big) \Big(2 A^{(1)} + F^{(0)} \Big) \Big(3 A^{(1)} + F^{(0)} \Big) \Big)$$

ACKNOWLEDGEMENT

We wish to thank Dr. S. K. A. Seyed Yagoobi for his careful reading the article and for his constructive comments.

References

- [1] A. V. TURBINER, *Commun. Math. Phys.***118**, 467 (1988).
- [2] M. A. SHIFMAN AND A. V. TURBINER, *Commun. Math. Phys.***120**, (1989), 347.
- [3] A. C. USHVERIDZE, *Sov. J. Part. Nucl.***20**504(1988).
- [4] M. A. SHIFMAN, *Int. J. Mod. Phys.***A 4** (1989),2897.
- [5] V. V. ULYANOV AND O. B. ZALAVSKII, *Phys. Rep.* **216**, 179 (1992).
- [6] C. M. BENDER AND G. V. DUNNE, *J. Math. Phys.* **37**(1996), 6.
- [7] M. A. JAFARIZADEH AND H. FAKHRI, *Annals of Physics* **262** (1998), 260.
- [8] M. A. JAFARIZADEH AND H. FAKHRI, *Physics Letters* **A230** (1997), 157.
- [9] Z. X. WANG AND D. R. GUO, “*Special Functions*”, World Scientific Publishing Co. LTD, (1989).
- [10] E. A. CODDINGTON AND N. LEVINSON, “*Theory of Ordinary Differential Equations*”, Tata McGraw-Hill Publishing Co. LTD, New Dehli, (1983).